

## AUTHOR INDEX

### Volume 40 (1988)

- |  |             |
|--|-------------|
| Bard, J.F., Convex two-level optimization  | (1) 15- 27  |
| Byrd, R.H., R.B. Schnabel and G.A. Shultz, Approximate solution of the trust region problem by minimization over two-dimensional subspaces | (3) 247-263 |
| Chaney, R.W., Second-order necessary conditions in semi-smooth optimization  | (1) 95-109  |
| Coleman, Th.F., A chordal preconditioner for large-scale optimization  | (3) 265-287 |
| Fiacco, A.V. and J. Kyparisis, Computable bounds on parametric solutions of convex problems  | (2) 213-221 |
| Fleischmann, B., A new class of cutting planes for the symmetric travelling salesman problem   | (3) 225-246 |
| Gay, D.M., Erratum   | (1) 111     |
| Glover, F. and D. Klingman, Layering strategies for creating exploitable structure in linear and integer programs                          | (2) 165-181 |
| Goldfarb, D. and S. Mehrotra, Relaxed variants of Karmarkar's algorithm for linear programs with unknown optimal objective value           | (2) 183-195 |
| Goldfarb, D. and S. Mehrotra, A relaxed revision of Karmarkar's method   | (3) 289-315 |
| Han, S.-P., A successive projection method   | (1) 1- 14   |
| Hoffman, A.J., On greedy algorithms for series parallel graphs   | (2) 197-207 |
| Imai, H., On the convexity of the multiplicative version of Karmarkar's potential function   | (1) 29- 32  |
| Klingman, D., <i>see</i> F. Glover   | (2) 165-181 |
| Kyparisis, J., <i>see</i> A.V. Fiacco  | (2) 213-221 |
| Mahjoub, A.R., On the stable set polytope of a series-parallel graph   | (1) 53- 57  |
| Mehrotra, S., <i>see</i> D. Goldfarb   | (2) 183-195 |
| Mehrotra, S., <i>see</i> D. Goldfarb   | (3) 289-315 |
| Meyer, G.G.L., Convergence of relaxation algorithms by averaging   | (2) 205-212 |
| Mifflin, R., Erratum   | (2) 223     |
| Pochet, Y. and L.A. Wolsey, Lot-size models with back-logging: Strong reformulations and cutting planes                                    | (3) 317-335 |
| Renegar, J., A polynomial-time algorithm, based on Newton's method, for linear programming   | (1) 59- 93  |
| Renegar, J. Rudiments of an average case complexity theory for piecewise-linear path following algorithms                                  | (2) 113-163 |
| Schnabel, R.B., <i>see</i> R.H. Byrd   | (3) 247-263 |
| Schultz, G.A., <i>see</i> R.H. Byrd  | (3) 247-263 |
| Tobin, R.L., A variable dimension solution approach for the general spatial price equilibrium problem                                      | (1) 33- 51  |
| Wolsey, L.A., <i>see</i> Y. Pochet   | (3) 317-335 |



